

Scenario 1:

You want to treat your milking goat herd with Safe-guard. You have 75 does that 200 pounds. You administer 4.6 ml of Safe-guard orally to each lactating doe. You record a 6 day slaughter withdrawal in the records.

Is this **GOOD**, **BAD**, or **ILLEGAL**?

LABEL INFORMATION:

- Used in goats for the removal and control of: Stomach worms (adults): *Haemonchus contortus* and *Teladorsagia circumcincta*

DIRECTIONS:

Determine the proper dose according to estimated body weight. Administer orally. The recommended dose of 2.3 mg/lb (5 mg/kg) of body weight is achieved when 2.3 mL of the drug are given for each 100 lbs body weight.

RESIDUE WARNINGS: Goats must not be slaughtered for food within 6 days following treatment. Because a withdrawal time in milk has not been established, do not use in lactating goats.

Scenario 2:

You want to treat your sheep with Ivomec drench. The sheep average 100 pounds. You administer 12 ml of Ivomec orally to each animal. You record an 11 day slaughter withdrawal in the records.

Is this **GOOD**, **BAD**, or **ILLEGAL**?

LABEL INFORMATION:

- IVOMEC Drench for Sheep provides treatment and control of adult and fourth-stage larvae of the following parasites: *Haemonchus contortus*, *Ostertagia circumcincta*, *Trichostrongylus axei*, *T. colubriformis*, *Cooperia curticei*, *Nematodirus spathiger*, *N. battus*, and *Oesophagostomum columbianum*; *Dictyocaulus filaria*; *Oestrus ovis*. *Haemonchus placei*, *Cooperia oncophora*, *Strongyloides papillosus*, *Oesophagostomum venulosum*, *Trichuris ovis*, and *Chabertia ovina*.

DIRECTIONS:

- IVOMEC Drench for Sheep is formulated to deliver orally at a volume of 3.0 mL per 26 lb body weight.

RESIDUE WARNING: Do not treat sheep within 11 days of slaughter.

Scenario 3:

You ran a fecal exam and determined whipworms (*Trichuris*) is the primary parasite on your farm. You want to treat your replacement heifers with Ivomec Plus. The heifers average 880 pounds. You administer 8 ml of Ivomec SQ to each animal. You record a 30 day slaughter withdrawal in the records.

Is this **GOOD**, **BAD**, or **ILLEGAL**?

LABEL INFORMATION:

- *Haemonchus*, *Ostertagia*, *Trichostrongylus*, *T. colubriformis*, *Cooperia*, *Nematodirus spathiger*, *N. battus*, and *Oesophagostomum c*; *Dictyocaulus filaria*; *Strongyloides*, *Oesophagostomum venulosum*, and *Chabertia ovina*, *Fasciola hepatica*, *Hypoderma bovis*

DIRECTIONS:

- IVOMEC Plus is formulated to deliver the recommended dose level of 200 mcg ivermectin/kg and 2 mg clorsulon/kg given subcutaneously behind the shoulder at the rate of 1 mL per 110 lb (50 kg) body weight.

RESIDUE WARNING: Do not treat cattle within 21 days of slaughter. Because a withdrawal time in milk has not been established, do not use in female dairy cattle of breeding age. A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

Scenario 4:

Parasites have been a continuous problem on your farm. Your vet recommends that you use Safe-guard to your lambs. You give each 50 pound lamb 1.2 ml orally and record a 30 day slaughter withdrawal in the records.

Is this **GOOD**, **BAD**, or **ILLEGAL**?

LABEL INFORMATION:

- Used in goats for the removal and control of: Stomach worms (adults): *Haemonchus contortus* and *Teladorsagia circumcincta*

DIRECTIONS:

Determine the proper dose according to estimated body weight. Administer orally. The recommended dose of 2.3 mg/lb (5 mg/kg) of body weight is achieved when 2.3 mL of the drug are given for each 100 lbs body weight.

RESIDUE WARNINGS: Goats must not be slaughtered for food within 6 days following treatment. Because a withdrawal time in milk has not been established, do not use in lactating goats.